



## Drought analysis in Mediterranean Region Akdeniz Bölgesinde kuraklık analizi

Ülker GÜNER BACANLI<sup>1\*</sup>, Gözde Nur AKŞAN<sup>2</sup>

<sup>1</sup>Department of Civil Engineering, Engineering Faculty, Pamukkale University, Denizli, Turkey.  
ugbacanli@pau.edu.tr, gozdeaksann@gmail.com

Received/Geliş Tarihi: 19.10.2018, Accepted/Kabul Tarihi: 24.01.2019  
\* Corresponding author/Yazışılan Yazar

doi: 10.5505/pajes.2019.64507  
Research Article/Araştırma Makalesi

### Abstract

Kuraklık gizlice gelişen bir doğal afettir. Bu çalışmada Standartlaştırılmış Yağış Evapotranspirasyon İndeksi (SPEI) Türkiye'nin Akdeniz Bölgesi'nde ilk kez uygulanmıştır. 8 meteoroloji gözlem istasyonunun sıcaklık ve yağış verileri kullanılmıştır. Verilere göre 8 istasyon (Adana, Antalya, Burdur, Hatay/Antakya, Isparta, Kahramanmaraş, Mersin ve Osmaniye) 1970-2018 yılları arasında gözlem yapılmıştır. Her bir istasyon için 1, 3, 6, 9 ve 12 aylık SPEI değerlerinin frekans analizleri hesaplanmıştır. Aylık (1, 3, 6, 9 ve 12 aylık) frekans değerleri arasında kurak ve sulak dönemlerin dağılımlarının karşılaştırılmasının yapılması amaçlanmıştır. Bulunan SPEI değerlerinin kuraklık sınıflarında ne kadar mevcut olduğu ve bu mevcudiyet üzerinden karşılaştırmalar yapılmıştır. Sonuç olarak Akdeniz Bölgesindeki tüm istasyonlarda elde edilen veriler hafif kuraklık ile normale yakınlık arasındadır. Hem normale yakın hem de kurak durumlarda Mersin maximum değerler almıştır. Minimum değerlere bakıldığında ise diğer istasyonlara kıyasla Adana hem sulak hem de kurak durumlarda en az yüzdellik değerlerine sahiptir.

**Keywords:** Kuraklık, Sıcaklık, Yağış, Standartlaştırılmış yağış evapotranspirasyon indeksi (SPEI)

### Öz

Drought is a natural disaster developing secretly. In this study, Standardized Precipitation Evapotranspiration Index (SPEI) has been applied for in Turkey's Mediterranean Region. Temperature and precipitation data were used for 8 meteorological observation stations. According to the data, 8 stations (Adana, Antalya, Burdur, Hatay/Antakya, Isparta, Kahramanmaraş, Mersin and Osmaniye) were observed between 1970-2018. Frequency analyzes of SPEI values of 1, 3, 6, 9 and 12 months were calculated for each station. Monthly (1, 3, 6, 9 and 12 months) frequency values between the distribution of ear and wetlands is intended to make a comparison. Comparisons were made on how long the SPEI values were found in drought classes and on this availability. As a result, the data obtained from all stations in the Mediterranean region are between mild dry and near to normal. Mersin has the maximum value both in near to normal and dry conditions. When the minimum values are considered, Adana has the least percentage values in both wetness and dryness conditions compared to other stations.

**Anahtar kelimeler:** Drought, Temperature, Precipitation, Standardized precipitation evapotranspiration index (SPEI)

## 1 Introduction

Drought is a world-wide effect of climate change and climatic events including values below average precipitation [1],[2]. It is a time-dependent phenomenon that is affected by many parameters [3]. The parameter used varies according to drought varieties. In this reason there is no specific formula for drought [3],[4]. In fact, it can be categorized as follows;

- Eteorological; it is defined as the precipitation falls significantly below normal values over a long period of time,
- Agricultural; it is a dry grounded period resulting from arid soils, low temperatures, more than expected precipitation events or more than normal evaporation,
- Hydrological; hydrological drought is associated with the effects of groundwater resources, surface waters or precipitation periods and,
- Socio-economic; the drought stage in which the social and economic impacts of water scarcity are felt prominently and the supply in the economy falls below demand due to drought [5]-[8].

Each natural hazard varies in various forms. According to this definition, forms of diversity are divided into three groups [9]. The first one, drought is a continuing phenomenon. When it starts, it will not be known but it will not end. Although the effects belong to a certain region, the whole world is under the influence of this phenomenon [10]. The second one, there is no precise definition of drought. The third one is the state in which the solution can be obtained because the drought is not a

sudden event. It can be controlled by appropriate monitoring and research [11]-[14].

As drought started to shape in a serious dimension in Africa, Alaska, Canada and Eurasia starting from 1950, scientists interested in this matter took notice and people started to find indices to minimize the drought [15]. Many indexes have been developed for the calculation of droughts. Some of these are Standardized Precipitation Index (SPI), Reclamation Drought Index (RDI), Effective Drought Index (EDI), Palmer Drought Severity Index (PDSI) and Standardized Precipitation Evapotranspiration Index (SPEI) [3],[16]-[19]. The purpose of all these indices is to detect and work on the arid regions that are now and in the future. It is necessary to investigate the detected regions and to minimize the risks that may occur.

Drought indices should include features such as determining drought and finding out how much the area is spreading. At the same time, they should not be same each other and be able to compare. SPEI uses both precipitation and temperature data to determine the region's drought. SPEI has emerged to prevent SPI problems. It is an advantageous drought index according to SPI [19].

The powerful feature of SPEI is that it allows for more accurate results with the help of numerical data. It is also affected by more than one factor, leading to more precise results [20]. Only precipitation and temperature values are sufficient for SPEI in climatic (meteorological) events. The results of time scales will be sufficient [21]. At the same time, we can say that SPEI covers the SPI because it considers the precipitation effect [1],[15]. In

fact, SPEI includes parameters used to obtain other drought indices too. However, the main advantage compared to other indices is that a multi-scalar character combines the capacity of evapotranspiration affected by temperature and drought severity, end and start time [7].

Reference Evapotranspiration (ET<sub>o</sub>) value is needed to calculate SPEI. ET<sub>o</sub> can be obtained in more than one way [15],[20],[22]-[24]. In addition to these methods, hybrid models (ARIMA-ANN, Wavelet-ANN (WANN) and WANFIS), log-linear models can be used to find these indices [14].

In this study, the main objective is to assess of meteorological droughts in the Mediterranean region of Turkey. Many drought indices have been used in the past to present day Turkey. Examples of these are the Standardized Precipitation Index (SPI), the Percentage of Normal Precipitation and the Palmer Drought Severity Index (PDSI). However, in this study, applied in the world of but which for in Turkey will be applied drought index SPEI it is used. Frequency analyzes of SPEI values of 1, 3, 6, 9 and 12 months were calculated for each station.

## 2 Methodology

### 2.1 Study area

There are 8 provinces of the Mediterranean Region in an area of 89.493 km<sup>2</sup>. In this study, between 1970-2018 it was aimed to calculate the drought for 8 meteorological stations in the Mediterranean Region (Figure 1).




Figure 1: Mediterranean region.

Table 1 provides data for the region (altitude, average rainfall, etc.). Using this data, the operations in the method section were performed.

Table 1: Data on the study area.

	Latitude (North)	Height (m)	Observation Time (year)	Average rainfall (mm/year)	Average temperature (mm)
Adana	37	23	49	656	19.2
Antalya	36.88	39	29	1119	18.6
Burdur	37.46	950	49	419	13.3
Hatay/ Antakya	36.4	85	49	1113	18.4
Isparta	37.76	1035	49	534	12.2
Kahramanmaraş	37.58	568	49	712	16.9
Mersin	36.8	6	49	588	19.4
Osmaniye	37.2	150	33	788	18.5

The evaluated monthly precipitation and temperature data were measured by the Turkish State Meteorological Services [DMI].

### 2.2 Methods

The SPEI is based on the Potential Evapotranspiration (PET) balance, which is the monthly climate balance. Precipitation and temperature values are taken into account when calculating the SPEI. This index can be calculated over several time periods. For this index benefit from the Table 2 created by [25]. Table 1 makes it easy to comment and monitor [26].

Table 2: Drought classification according to SPEI categories based on [25].

Categories	
Exceptionally wet	$\geq +2$
Severely wet	$\geq +1.5$ to $< +2$
Mild wet	$\geq +1$ to $< +1.5$
Near to Normal	$> -1$ to $< +1$
Mild dry	$> -1.5$ to $\leq -1$
Severely dry	$> -2$ to $\leq -1.5$
Exceptionally dry	$\leq -2$

The following operations must be performed to obtain SPEI. The following steps must be taken to obtain SPEI. First of all, we start with PET calculation. PET calculation can be calculated with more than one method. For example, Thornwaite, Blaney-Cridle etc. mentioned. However, Thornwaite (1948) method was preferred because it could be calculated more easily in this study.

$$PET = 16K \left( \frac{10 * T_{mm}}{I} \right)^m \quad (1)$$

$K$  is a function calculated by the correction coefficient of the latitude and month,  $T_{mm}$  is monthly-mean temperature (°C),  $I$  is a heat index and  $m$  is a coefficient depending upon  $I$ .

Monthly precipitation and  $PET$  values are used in millimeters units.

$$D_i = P_i - PET_i \quad (2)$$

$P$  and  $PET$  is calculated for the month  $i$ . This difference shows a simple meteorological water balance [27].

$$D_n^k = \sum_{i=0}^{k-1} (P_{n-i} - PET_{n-i}), \quad n \geq k \quad (3)$$

For the different  $D$  series,  $k$  (month) is the time scale of the cluster and  $n$  is the calculation number. The  $D$  values are undefined for  $k > n$ .

The PWMs (probability-weighted moments) of order  $s$  are calculated as,

$$w_s = \frac{1}{N} \sum_{i=1}^N (1 - F_i)^s * D_i \quad (4)$$

The resulting  $w_s$  values are used to find  $\alpha$ ,  $\beta$  and  $\gamma$ .

$F_i$  is a frequency estimator,  $N$  is the number of data points. The probability distribution of series  $D$  is as follows;

$$F(x) = \left[ 1 + \left( \frac{\alpha}{x - \gamma} \right)^\beta \right]^{-1} \quad (5)$$

$F(x)$  formula,  $\alpha$ ,  $\beta$  and  $\gamma$  contain scale, shape and origin parameters [28]. For the  $D$  range,  $\gamma > D < \infty$ .

As given in Equation 6, several scientists have suggested some approaches to the calculation of SPEI [29];

$$SPEI = W - \frac{C_0 + C_1 * W + C_2 * W^2}{1 + d_1 * W + d_2 * W^2 + d_3 * W^3} \quad (6)$$

Where,

$$W = \sqrt{-2 \ln(P)} \quad (7)$$

$P$  is the probability of wear of a D value,  $P = 1 - F(x)$ . If  $P > 0.5$ , then then  $P$  is replaced by  $1-P$ . Changes the SPEI value sign obtained.  $W$  is a tool used to obtain  $P$ . Constants used for SPEI;  $C_0 = 2.515517$ ,  $C_1 = 0.802853$ ,  $C_2 = 0.010328$ ,  $d_1 = 1.432788$ ,  $d_2 = 0.189269$ ,  $d_3 = 0.001308$

The cumulative probability for time scales is calculated. The SPEI value is then obtained by converting the standard normal distribution to zero and to a variance [30],[31].

### 3 Results and discussions

The SPEI values were estimated on 1, 3, 6, 9- and 12-months' time scale conditions for all stations. Thornthwaite, Penman-Monteith and Hargreaves methods can be used to calculate the SPEI. However, in this study, we decided to use the Thornthwaite method because we understood that these methods were simple and useful. As example, it was seen SPEI values graphs for Adana from Figure 2, dry and wet season periods are observed to increase from 1 to 12 months.




Figure 2: 1, 3, 6, 9 and 12-month SPEI values for Adana station.

SPEI relative frequency results for all stations are given on Table 3-10.

The observed relative frequency of 1-month SPEI values for Adana was found to be near to normal and 67.47% maximum than other drought periods (Table 3). It was observed that the total of the data which is near to normal in the data of 6 months frequency is 52.94% minimum than the other drought

varieties. Exceptionally dry period was observed in all time periods but not in other periods. However, throughout the arid interval, the frequency of 1-month SPEI viewing was generally lower than other frequencies. As it can be seen from Table 3, 1-month frequency values near to normal and exceptionally wet, 3, 6, and 9 months frequency values are observed in mild dry or mild wet intervals and 12-month frequency values are near to normal and exceptionally dry.

Table 3: Relative frequency percentage in Adana.

<b>ADANA</b>	1	3	6	9	12
Exceptionally wet	1.73	1.90	0.87	1.90	1.73
Severely wet	7.27	3.29	4.15	3.11	3.98
Mild wet	8.48	11.07	15.40	11.42	9.34
Near to Normal	67.47	60.03	52.94	60.21	63.32
Mild dry	9.52	21.28	24.57	20.59	14.88
Severely dry	3.63	2.42	2.08	2.77	6.75
Exceptionally dry	1.90	0.00	0.00	0.00	0.00

Table 4: Relative frequency percentage in Antalya.

<b>ANTALYA</b>	1	3	6	9	12
Exceptionally wet	2.30	2.01	1.44	2.59	2.87
Severely wet	5.17	4.60	5.75	4.02	3.16
Mild wet	8.05	10.63	13.22	11.49	12.07
Near to Normal	68.10	61.21	58.05	59.77	60.63
Mild dry	11.49	20.40	20.40	20.11	19.25
Severely dry	2.87	1.15	1.15	2.01	2.01
Exceptionally dry	2.01	0.00	0.00	0.00	0.00

Table 5: Relative frequency percentage in Burdur.

<b>BURDUR</b>	1	3	6	9	12
Exceptionally wet	1.90	0.87	1.04	1.21	1.38
Severely wet	4.84	3.63	3.11	3.98	3.81
Mild wet	10.55	13.32	15.74	11.94	11.59
Near to Normal	65.92	58.48	54.15	59.69	60.73
Mild dry	11.07	17.13	22.84	16.78	13.67
Severely dry	4.84	6.57	2.94	6.40	8.82
Exceptionally dry	0.87	0.00	0.17	0.00	0.00

Table 6: Relative frequency percentage in Hatay/Antakya.

<b>HATAY/ANTAKYA</b>	1	3	6	9	12
Exceptionally wet	1.90	1.21	1.04	1.21	1.21
Severely wet	5.88	4.84	5.02	4.84	5.19
Mild wet	9.86	12.11	13.84	11.76	11.07
Near to Normal	66.61	56.40	51.38	56.06	57.61
Mild dry	10.03	24.39	26.82	24.91	23.18
Severely dry	4.15	1.04	1.90	1.21	1.73
Exceptionally dry	1.56	0.00	0.00	0.00	0.00

Table 7: Relative frequency percentage in Isparta.

<b>ISPARTA</b>	1	3	6	9	12
Exceptionally wet	2.25	1.73	1.04	1.73	1.73
Severely wet	5.70	4.49	5.70	3.97	4.32
Mild wet	9.33	10.02	11.57	10.36	9.67
Near to Normal	66.15	61.14	57.51	60.97	62.52
Mild dry	11.40	15.54	20.90	16.06	13.99
Severely dry	3.97	7.08	2.25	6.91	7.77
Exceptionally dry	1.21	0.00	1.04	0.00	0.00

Table 8: Relative frequency percentage in Kahramanmaras.

<b>KAHRAMANMARAS</b>	1	3	6	9	12
Exceptionally wet	1.04	1.21	0.35	0.87	0.87
Severely wet	7.27	4.84	5.54	5.54	5.36
Mild wet	10.21	11.94	16.78	11.94	11.07
Near to Normal	65.22	57.79	51.73	56.92	58.82
Mild dry	9.52	21.97	24.57	22.32	20.42
Severely dry	5.71	2.25	1.04	2.42	3.46
Exceptionally dry	1.04	0.00	0.00	0.00	0.00

Table 9: Relative frequency percentage in Mersin.

<b>MERSİN</b>	1	3	6	9	12
Exceptionally wet	2.25	1.21	1.21	1.21	1.38
Severely wet	4.84	3.81	4.15	5.02	3.63
Mild wet	9.00	14.36	16.61	12.63	12.11
Near to Normal	68.17	59.34	55.19	59.69	63.32
Mild dry	9.17	17.47	21.45	17.30	13.49
Severely dry	5.54	3.81	1.21	4.15	6.06
Exceptionally dry	1.04	0.00	0.17	0.00	0.00

Table 10: Relative frequency percentage in Osmaniye.

<b>OSMANIYE</b>	1	3	6	9	12
Exceptionally wet	3.14	2.62	2.88	2.36	2.09
Severely wet	2.62	4.19	4.45	4.19	4.97
Mild wet	10.21	9.16	8.64	9.42	8.64
Near to Normal	62.83	67.80	65.45	67.54	68.06
Mild dry	14.14	7.33	9.16	7.07	7.07
Severely dry	6.81	6.81	8.90	8.12	7.33
Exceptionally dry	0.26	2.09	0.52	1.31	1.83

Antalya can be interpreted as taking place in the normal drought class. Near to normal, a maximum of 68.10 percent is available, and a minimum of 58.05 percent is obtained. Drought for Antalya has a minimum value of 16.37% and maximum value of 22.12%. Minimum 15.52% and maximum 20.40% values were obtained in wetness (Table 4). When the comparison is made according to Table 4, it is seen that 1-month frequency values are near to normal and exceptionally dry, 3-month frequency values are near to normal and mild dry, 6 and 9-month frequency values are mild dry, or mild wet and 12-month frequency values are exceptionally wet.

It can be said that the dryness for Burdur is between 25.95-16.78%. The wetness was observed to be between 19.9-16.78% (Table 5). However, it is concluded that the station is in near to normal condition. In the near to normal drought class, it is seen that 1-month frequency has a maximum value of 65.92%. According to Table 5, frequency values of 1 month were near to normal, frequencies of 3, 6 and 9 months were mild dry or mild wet and 12 months frequency was found to be severely dry.

According to all frequencies in Hatay / Antakya, dryness varies between 15.74% and 28.72%. The wetness is between 17.47% and 19.90% (Table 6). But the station near to normal has the highest percentage. This percentage is 66.61% in the 1-month frequency. It has been concluded that all frequencies in normal drought class exceed 50%. When compared with Table 6, it is seen that the frequency values of 1 month, 3, 6 and 9 months of the frequency values of 1 month were mild dry or mild wet and 12 months of frequency values of -1.5 < to <+1.5. Mild dry is very low compared to other frequencies in the 1-month frequency.

For Isparta, results were found between 66.15% and 57.51% in the near to normal drought class. However, if the Table 7 was placed in a general class, it was concluded that the dryness was

between 16.58% and 24.18% and the wetness was between 15.72% and 18.31% (Table 7). As It can be seen from Table 7, 1-month of near to normal frequency values, 3, 6 and 9 months were mild dry or mild wet, 12 months frequency values were close to near to normal and severely dry. When the Table 8 for Kahramanmaras is examined, it is concluded that the near to normal drought class has the highest percentage. Near to normal drought class has a minimum value with a 1-month frequency value (65.22%) and a maximum 6-month frequency value (51.73%). In addition, dryness and wetness of the station were divided into two, dryness between 16.26-25.61%, wetness resulted in values of 17.30-22.66% (Table 8). As It can be seen from Table 8, 1-month frequency values of near to normal and severely wet, 3-month and 6-month and 9-month frequency values of mild dry or mild wet and 12-month frequency values of near to normal and mild dry conditions were observed.

The maximum value for Mersin station was found in the near to normal drought class at 1-month frequency (68.17%). When the Table 9 is classified as dry and wet, the dry class takes a minimum of 15.74% and the maximum is 22.84%. The wet class was between 16.09% and 21.97% (Table 9). As it can be seen from Table 9, frequency values of 1 month are near to normal and frequencies of 3, 6 and 9 months are mild dry, or mild wet and 12-month frequency values are near to normal or severely dry. When the Table 10 for Osmaniye was examined, the maximum value in the near to normal drought class was found at the frequency of 12 months (68.06%). The minimum value was obtained at 1-month frequency (62.83%). The near to normal drought class is the highest in all frequencies. In addition, all frequencies in the near to normal drought class exceed 60%. Unlike other stations, the wetness at this station has the same percentage (15.97%) for frequencies 1, 3, 6, 9.

Dryness also varies between 16.23% and 21.20% (Table 10). And unlike other illusions, the frequency values of 1 month were mild dry or mild wet, the frequency values of 3 and 9 months were near to normal, the frequency values of 6 months were exceptionally dry and 12 months frequency values were near to normal. Mediterranean Anatolia Region are the regions most affected by arid conditions in early 1970s and early 1990s. If we accept Antalya as the starting point, the drought towards the northwest is turning to the previous year. Drought towards the northeast is also over the previous year. In other words, the same chart is displayed on the right side in both directions. Drought last for 3 years (Figure 3).

#### 4 Conclusions

Drought is a natural disaster that causes significant problems in life. Drought analysis and management is very important in combating drought problems. Because drought is a very complex phenomenon, each drought is characterized by different properties. SPEI were used to determine drought and effective drought management in many countries [20]. The SPEI was applied for in Turkey by this study. In the Mediterranean region between 1970 and 2018, mild, severely and exceptionally levels of drought were observed.

The remarkable drought interval at all stations is  $> -1$  to  $< +1$ . In other words, the country is located in the near-normal value range. In general, the near to normal drought percentage of the 1-month frequency is higher than the percentage of the 12-month frequency.

It was seen that dryness varies between 15.05% and 28.72% all stations. The wetness varies between 15.05% and 22.66%. Looking at the region in general, it can be concluded that the near to normal drought class has the highest percentages. Mersin has the maximum percentage with a 1-month frequency value (68.17%) in normal drought class. Hatay/Antakya has a minimum percentage with a frequency of 6 months (51.38%).

The Mediterranean region may also face the danger of agricultural and hydrological drought seen later than

meteorological drought. Drought prevention plan can be created which to reduce the potential drought effects of the Mediterranean region. In addition, the use of water resources can be regulated.

#### 5 References

- [1] Kabat P, Schulze RE, Hellmuth ME, Veraart JA. Coping with Impacts of Climate Variability and Climate Change in Water Management: A Scoping Paper. International Secretariat of the Dialogue on Water and Climate, Wageningen, Netherlands, 2003.
- [2] Trenberth KE, Dai A, Van der Schrier G, Jones PD, Barichivich J, Briffa KR and Sheffield J. "Global warming and changes in drought". *Natural Climate Change*, 4(1), 17-22, 2014.
- [3] Potop V, Možný M, Soukup J. "Drought evolution at various time scales in the lowland regions and their impact on vegetable crops in the Czech Republic". *Agricultural and Forest Meteorology*, 156, 121-133, 2012.
- [4] Keyantash J, Dracup JA. "The quantification of drought: An evaluation of drought indices". *Bulletin of the American Meteorological Society*, 83, 1167-1180, 2002.
- [5] Wilhite DA, Glantz MH. "Understanding: the drought phenomenon: The role of definitions". *Water International*, 10(3), 111-120, 1985.
- [6] Mouatadid S, Raj N, Deo RC, Adamowski JF. "Input selection and data-driven model performance optimization to predict the standardized precipitation and evaporation index in a drought-prone region". *Atmospheric Research*, 212, 130-149, 2018.
- [7] Dai A. "Drought under global warming: a review". *Wiley Interdisciplinary Reviews: Climate Change*, 2(1), 45-65, 2011.
- [8] Potop V, Možný M, Boroneanț C, Štěpánek P and Skalák P. "Observed spatiotemporal characteristics of drought on various time scales over the Czech Republic". *Theoretical and Applied Climatology*, 115(3-4), 563-581, 2014.



Figure 3: Years where the region has the highest drought in all frequencies.

- [9] Wilhite DA, Sivakumar MVK, Wood DA. "Warning systems for drought preparedness and drought management". *World Meteorological Organization*, Lisbon, Portugal, 5-7 September 2000.
- [10] Tannehill IR. *Drought its Causes and Effects*. New Jersey, USA, Princeton University, 1947.
- [11] Alam NM, Adhikary PP, Jana C, Kaushal R, Sharma NK, Avasthe RK, Ranjan R, Mishra PK. "Application of Markov model and standardized precipitation index for analysis of droughts in Bundelkhand region of India". *Journal of Tree Sciences*, 31(1&2), 46-53, 2012.
- [12] Alam NM, Mishra PK, Jana C, Adhikary PP. "Stochastic model for drought forecasting for Bundelkhand region in Central India". *Indian Journal of Agricultural Sciences*, 84(2), 79-84, 2014.
- [13] Alam NM, Ranjan R, Adhikary PP, Kumar A, Jana C, Panwar S, Mishra PK, Sharma NK. "Statistical modeling of weekly rainfall data for crop planning in Bundelkhand region of Central India". *India Section B: Biological Sciences*, 44(3), 336-342, 2016.
- [14] Alam NM, Sharma GC, Moreira E, Jana C, Mishra PK, Sharma NK, Mandal D. "Evaluation of drought using SPEI drought class transitions and log-linear models for different agro-ecological regions of India". *Physics and Chemistry of the Earth*, 100, 31-43, 2017.
- [15] Chen H, Sun J. "Changes in drought characteristics over china using the standardized precipitation evapotranspiration index". *Journal of Climate*, 28, 5430-5447, 2015.
- [16] Bamimahd SM, Khalili D. "Factors Influencing Markov Chains Predictability Characteristics, Utilizing SPI, RDI, EDI and SPEI Drought Indices in Different Climatic Zones". *Water Resources Management*, 27, 3911-3928, 2013.
- [17] Wang W, Zhu Y, Xu R, Liu J. "Drought severity change in China during 1961-2012 indicated by SPI and SPEI". *Natural Hazards*, 75(3), 2437-2451, 2014.
- [18] Hou M, Li H, Zou X, An W, Gao G, Zhao H. "Timescale differences between SC-PDSI and SPEI for drought monitoring in China". *Physics and Chemistry of the Earth*, 102, 48-58, 2017.
- [19] Stagge JH, Tallaksen LM, Gudmundsson L, Van Loon A and Stahl K. "Candidate Distributions for Climatological Drought Indices (SPI and SPEI)". *International Journal of Climatology*, 35(13), 4027-4040, 2015.
- [20] van der Schrier G, Barichivich J, Briffa KR, Jones PD. "A scPDSI-based global data set of dry and wet spells for 1901-2009". *Journal of Geophysical Research: Atmospheres*, 118(10), 4025-4048, 2013.
- [21] Huang YF, Soh YW, Koo CH, Fung KF. "Application of artificial intelligence models for the prediction of standardized precipitation evapotranspiration index (SPEI) at Langat River Basin, Malaysia". *Computers and Electronics in Agriculture*, 144, 164-173, 2018.
- [22] Donohue RJ, McVicar T, Roderick ML. "Assessing the ability of potential evaporation formulations to capture the dynamics in evaporative demand within a changing climate". *Journal of Hydrology*, 386(1-4), 186-197, 2010.
- [23] Vicente-Serrano SM, Beguería S, Azorin-Molina C, Schrier Van der G. "Contribution of precipitation and reference evapotranspiration to drought indices under different climates". *Journal of Hydrology*, 526, 42-54, 2015.
- [24] Dai A. "Increasing drought under global warming in observations and models". *Natural Climate Change*, 3(1), 52, 2013.
- [25] McKee TB, Doesken NJ, Kleist J. "The relationship of drought relative frequency and duration to time scales". *8<sup>th</sup> Conference on Applied Climatology*, California, USA, 17-22 January 1993.
- [26] Abramowitz M, Stegun IA. *Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables*. New York, USA, Dover Publications, 1965.
- [27] Hernandez EA, Uddameri V. "Standardized precipitation evaporation index (SPEI)-based drought assessment in semi-arid south Texas". *Environmental Earth Sciences*, 71(6), 2491-2501, 2014.
- [28] Mouatadida S, Rajb N, Deob RC, Adamowskic JF. "Input selection and data-driven model performance optimization to predict the Standardized Precipitation and Evaporation Index in a drought-prone region". *Atmospheric Research*, 212, 130-149, 2018.
- [29] Edwards DC, McKee TB. "Characteristics of 20<sup>th</sup> century drought in the United States at multiple time scales". *Atmospheric Science Thesis*, 634, 1-30, 1997.
- [30] Onusluel Gul G. and Kuzucu A. "Analysis of drought severity in Seyhan river basin". *European Water*, 60, 211-217, 2017.
- [31] Vicente-Serrano SM, Beguería S, Lopez-Moreno JL. "A Multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index". *Journal of Climate*, 23(7), 1696-1718, 2010.