



On Modal Operators over the Generalized Intuitionistic Fuzzy Set

Ezzatallah BALOUI JAMKHANEH^{1,*}, Saralees NADARAJAH²

¹Department of Statistics, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran

²School of Mathematics, University of Manchester, Manchester, UK

Article Info

Received: 16/06/2017

Accepted: 23/12/2017

Keywords

Generalized
intuitionistic fuzzy set
Intuitionistic fuzzy set
Level operator
Modal operator

Abstract

In recent years, different operators (modal, topological, level, negation and other types) have been defined over intuitionistic fuzzy sets. In this note, modal types of operators over an extended generalized intuitionistic fuzzy set GIFSB are proposed. Some of the basic properties of the new operators are derived.

1. INTRODUCTION

In the real world, we frequently deal with vague, imprecise or insufficient information. Fuzzy sets (FSs) were introduced to deal with this reality. In the fuzzy sets theory, membership and non- membership degrees are complementary, i.e., the sum of both degrees of an element belonging in a fuzzy set is 1. However, there are situations where the two degrees are not complementary, mainly because of hesitation. In an intuitionistic fuzzy set (IFS), each element is assigned membership and non-membership degrees, where the sum of the two degrees does not exceed 1. Accordingly, an IFS is more applicable than a FS for representing vague information since it considers a degree of hesitation. IFSs and their generalizations can be useful in problems such as decision making problems, sales analysis, new product marketing, and financial services. The modal operators have been known to be important tools for IFSs and allow for a more detailed estimation of information.

IFSs were introduced by Atanassov [1]. Since then different types of operators have been defined over IFSs. They can be classified into several groups: modal operators, topological operators, level operators, negation operators and other types. Atanassov [2] defined several operators for the theory of IFSs. He discussed the relations between the classical negation operator and the two standard modal operators “necessity” and “possibility”. Atanassov [1,3-4] defined topological operators over IFSs, and derived their basic properties. Atanassov [5] considered modal operators defined over IFSs. Cornelis et al. [6] considered extended modal operators as tools for constructing inclusion indicators over IFSs. Atanassov and Dimitrov [7] introduced different forms of negation operators. Hinde and Atanassov [8] studied some relations between intuitionistic fuzzy negations and intuitionistic fuzzy modal operators. Parvathi and Geetha [9] defined some level operators, max-min implication operators and $P_{\alpha,\beta}$, $Q_{\alpha,\beta}$ operators on temporal IFSs. Atanassov [10] introduced two new operators that partially extend the intuitionistic fuzzy operators from modal type. Srinivasan and Palaniappan [11] defined some operators and established their properties over IFSs of root type. Sharma [12] studied the impact of the modal operator $F_{\alpha,\beta}(A)$ on

*Corresponding author, e-mail: e_baloui2008@yahoo.com

intuitionistic fuzzy groups and proved that many properties of intuitionistic fuzzy subgroups like normality, commutativity and cyclic groups remain invariant under the modal operator. Yilmaz and Cuvalcioglu [13] introduced $N_B(A)$ and $N_{B^*}(A)$ operators on temporal IFSs and examined their properties. Baloui Jamkhaneh and Nadarajah [14] introduced a new generalized IFS ($GIFS_B$) and some operators over $GIFS_B$. Baloui Jamkhaneh [15] and Baloui Jamkhaneh and Nadi Ghara [16] defined level operators over the $GIFS_B$ s. In 2017 Baloui Jamkhaneh and Garg [17] considered some new operations over the generalized intuitionistic fuzzy sets and their application to decision making process. All operations defined over IFSs were transformed to the $GIFS_B$ case.

The aim of this note is to define different modal type operators over $GIFS_B$ and to derive their properties. This study shows the exibility of $GIFS_B$ s in terms of different operators. The study is expected to encourage applicability of $GIFS_B$ to the many areas of application of IFSs like sustainable energy planning, image fusion, agricultural production planning, medical diagnosis, pattern recognition, reservoir flood control operation, reliability optimization of complex systems, fault diagnosis and prediction of the best quality of two-wheelers, see Baloui Jamkhaneh and Nadarajah [14] for references and other areas of application of IFSs. The contents of this note are organized as follows: Section 2 states some definitions including the definition of $GIFS_B$ given in Baloui Jamkhaneh and Nadarajah [14]; Section 3 summarizes some results on $GIFS_B$ given in Baloui Jamkhaneh and Nadarajah [14]; Section 4 defines operators acting on $GIFS_B$ and derives their properties; some conclusions and future work are noted in Section 5.

2. BASIC CONCEPTS

Definition 2.1. [1] Let X be a non-empty set. An IFS A in X is defined as an object of the form $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$ where the functions $\mu_A : X \rightarrow [0,1]$ and $\nu_A : X \rightarrow [0,1]$ denote the degree of membership and non-membership functions of A respectively and $0 \leq \mu_A(x) + \nu_A(x) \leq 1$ for each $x \in X$.

Definition 2.2. [14] Let X be a non empty set. Generalized intuitionistic fuzzy sets A in X , is defined as an object of the form $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$ where the functions $\mu_A : X \rightarrow [0,1]$ and $\nu_A : X \rightarrow [0,1]$, denote the degree of membership and degree of non-membership functions of A respectively, and $0 \leq \mu_A(x)^\delta + \nu_A(x)^\delta \leq 1$ for each $x \in X$, and given fixed δ . The collection of all generalized IFSs is denoted by $GIFS_B(\delta, X)$.

Definition 2.3. [14] Let X be a non-empty set. Let A and B be two $GIFS_B$ s such that

$$A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}, \quad B = \{ \langle x, \mu_B(x), \nu_B(x) \rangle : x \in X \}.$$

Define the following operations on A and B :

- i. $A \subset B$ if and only if $\mu_A(x) \leq \mu_B(x)$ and $\nu_A(x) \geq \nu_B(x)$, $\forall x \in X$,
- ii. $A = B$ if and only if $\mu_A(x) = \mu_B(x)$ and $\nu_A(x) = \nu_B(x)$, $\forall x \in X$,
- iii. $A \cup B = \{ \langle x, \max(\mu_A(x), \mu_B(x)), \min(\nu_A(x), \nu_B(x)) \rangle : x \in X \}$,
- iv. $A \cap B = \{ \langle x, \min(\mu_A(x), \mu_B(x)), \max(\nu_A(x), \nu_B(x)) \rangle : x \in X \}$,
- v. $A + B = \{ \langle x, \mu_A(x)^\delta + \mu_B(x)^\delta - \mu_A(x)^\delta \mu_B(x)^\delta, \nu_A(x)^\delta \nu_B(x)^\delta \rangle : x \in X \}$,
- vi. $A \cdot B = \{ \langle x, \mu_A(x)^\delta \mu_B(x)^\delta, \nu_A(x)^\delta + \nu_B(x)^\delta - \nu_A(x)^\delta \nu_B(x)^\delta \rangle : x \in X \}$,
- vii. $\bar{A} = \{ \langle x, \nu_A(x), \mu_A(x) \rangle : x \in X \}$.

Definition 2.4. The degree of non-determinacy (uncertainty) of an element $x \in X$ to the $GIFS_B$ A is defined by $\pi_A(x) = (1 - \mu_A(x)^\delta - \nu_A(x)^\delta)^{\frac{1}{\delta}}$.

It can be easily shown that $\pi_A(x)^\delta + \mu_A(x)^\delta + \nu_A(x)^\delta = 1$.

3. REMARKS ON THE GENERALIZED INTUITIONISTIC FUZZY SETS

Let X is a non-empty finite set, $\alpha, \beta \in [0,1]$ and $A \in \text{GIFS}_B$, as $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$. Baloui Jamkhaneh and Nadarajah [14] introduced the following operators over GIFS_B and investigated some their properties:

- i. $\Box A = \{ \langle x, \mu_A(x), (1 - \mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$, (modal logic: the necessity measure),
- ii. $\Diamond A = \{ \langle x, (1 - \nu_A(x)^\delta)^{\frac{1}{\delta}}, \nu_A(x) \rangle : x \in X \}$, (modal logic: the possibility measure),
- iii. $C(A) = \{ \langle x, K, L \rangle : x \in X \}$, $K = \max_{y \in X} \mu_A(y)$, $L = \min_{y \in X} \nu_A(y)$, (topological: closure),
- iv. $I(A) = \{ \langle x, k, l \rangle : x \in X \}$, $k = \min_{y \in X} \mu_A(y)$, $l = \max_{y \in X} \nu_A(y)$, (topological: intersection),
- v. $D_\alpha(A) = \{ \langle x, (\mu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\nu_A(x)^\delta + (1 - \alpha) \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$,
- vi. $F_{\alpha, \beta}(A) = \{ \langle x, (\mu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\nu_A(x)^\delta + \beta \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$, $\alpha + \beta \leq 1$,
- vii. $G_{\alpha, \beta}(A) = \{ \langle x, \alpha^{\frac{1}{\delta}} \mu_A(x), \beta^{\frac{1}{\delta}} \nu_A(x) \rangle : x \in X \}$.

Theorem 3.1. For every GIFS_B A , we have

- i. $\Box \Box A = \Box A$,
- ii. $\Diamond \Diamond A = \Diamond A$,
- iii. $\Box \Diamond A = \Diamond A$,
- iv. $\Diamond \Box A = \Box A$,
- v. $D_\alpha(\Box A) = F_{\alpha, \beta}(\Box A) = \Box A$,
- vi. $D_\alpha(\Diamond A) = F_{\alpha, \beta}(\Diamond A) = \Diamond A$,
- vii. $D_\alpha(\bar{A}) = \overline{D_{1-\alpha}(A)}$,
- viii. $G_{\alpha, \beta}(A \cap B) = G_{\alpha, \beta}(A) \cap G_{\alpha, \beta}(B)$,
- ix. $G_{\alpha, \beta}(A \cup B) = G_{\alpha, \beta}(A) \cup G_{\alpha, \beta}(B)$,
- x. $\pi_{D_\alpha(A)}(x) = 0$,
- xi. $F_{\alpha, \beta}(D_\alpha(A)) = D_\alpha(A)$.

Proof. Proof is obvious.

4. MAIN RESULTS

Here, we will introduce new operators over the GIFS_B , which extend some operators in the research literature related to IFSs.

Definition 4.1. Letting $\alpha, \beta \in [0,1]$ and $A \in \text{GIFS}_B$, we define the operators of as follows

- i. $d_\alpha(A) = \{ \langle x, (\nu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + (1 - \alpha) \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$,
- ii. $f_{\alpha, \beta}(A) = \{ \langle x, (\nu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$, where $\alpha + \beta \leq 1$,
- iii. $g_{\alpha, \beta}(A) = \{ \langle x, \alpha^{\frac{1}{\delta}} \nu_A(x), \beta^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \}$,
- iv. $J_{\alpha, \beta}(A) = \{ \langle x, (\mu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta \nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$,
- v. $j_{\alpha, \beta}(A) = \{ \langle x, (\nu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta \mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$,
- vi. $H_{\alpha, \beta}(A) = \{ \langle x, (\alpha \mu_A(x)^\delta)^{\frac{1}{\delta}}, (\nu_A(x)^\delta + \beta \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$,
- vii. $h_{\alpha, \beta}(A) = \{ \langle x, (\alpha \nu_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \}$.

Theorem 4.1. For every $A \in GIFS_B$ and for every $\alpha, \beta \in [0,1]$, we have

- i. $d_\alpha(A) \in GIFS_B$,
- ii. $\alpha \leq \beta \Rightarrow d_\alpha(A) \subset d_\beta(A)$,
- iii. $d_0(A) = \overline{\diamond A}$,
- iv. $d_1(A) = \overline{\square A}$,
- v. $d_\alpha(\square A) = \overline{\square A}$,
- vi. $d_\alpha(\diamond A) = \overline{\diamond A}$.

Proof. Proofs (i) and (ii) are obvious.

(iii) Note that

$$\begin{aligned} d_0(A) &= \left\{ \langle x, (v_A(x)^\delta + 0\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + (1-0)\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}, \\ &= \left\{ \langle x, v_A(x), (\mu_A(x)^\delta + \pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}. \end{aligned}$$

Since $\pi_A(x)^\delta = 1 - \mu_A(x)^\delta - v_A(x)^\delta$ we have

$$d_0(A) = \left\{ \langle x, v_A(x), (1 - v_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\} = \overline{\diamond A}.$$

The proof is complete.

(iv) Follows by noting that

$$\begin{aligned} d_1(A) &= \left\{ \langle x, (v_A(x)^\delta + 1\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + (1-1)\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}, \\ &= \left\{ \langle x, (v_A(x)^\delta + \pi_A(x)^\delta)^{\frac{1}{\delta}}, \mu_A(x) \rangle : x \in X \right\}, \\ &= \left\{ \langle x, (1 - \mu_A(x)^\delta)^{\frac{1}{\delta}}, \mu_A(x) \rangle : x \in X \right\} = \overline{\square A}. \end{aligned}$$

(v) Note that

$$d_\alpha(\square A) = \left\{ \langle x, (v_{\square A}(x)^\delta + \alpha\pi_{\square A}(x)^\delta)^{\frac{1}{\delta}}, (\mu_{\square A}(x)^\delta + (1-\alpha)\pi_{\square A}(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

Since $\pi_{\square A}(x) = 0$, we have

$$d_\alpha(\square A) = \{ \langle x, v_{\square A}(x), \mu_{\square A} \rangle : x \in X \} = \overline{\square A}.$$

The proof is complete. The proof of (vi) is similar to that of (v).

Theorem 4.2. For every $A \in GIFS_B$ and for every $\alpha, \beta \in [0,1]$, where $\alpha + \beta \leq 1$, we have

- i. $f_{\alpha,\beta}(A) \in GIFS_B$,
- ii. $0 \leq \gamma \leq \alpha \Rightarrow f_{\gamma,\beta}(A) \subset f_{\alpha,\beta}(A)$,
- iii. $0 \leq \gamma \leq \beta \Rightarrow f_{\alpha,\beta}(A) \subset f_{\alpha,\gamma}(A)$,
- iv. $d_\alpha(A) = f_{\alpha,1-\alpha}(A)$,
- v. $f_{0,1}(A) = \overline{\diamond A}$,
- vi. $f_{1,0}(A) = \overline{\square A}$,
- vii. $\overline{f_{\alpha,\beta}(A)} = F_{\alpha,\beta}(\bar{A})$,
- viii. $\overline{f_{\alpha,\beta}(\bar{A})} = f_{\beta,\alpha}(A)$,
- ix. $f_{0,0}(A) = \bar{A}$,

- x. $f_{\alpha,\beta}(\Box A) = \overline{\Box A}$,
- xi. $f_{\alpha,\beta}(\Diamond A) = \overline{\Diamond A}$.

Proof. (i) Follows by noting that

$$\begin{aligned} \mu_{f_{\alpha,\beta}(A)}(x)^\delta + \nu_{f_{\alpha,\beta}(A)}(x)^\delta &= [(\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}]^\delta + [(\mu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}}]^\delta, \\ &= \nu_A(x)^\delta + \mu_A(x)^\delta + \pi_A(x)^\delta(\alpha + \beta), \\ &\leq \mu_A(x)^\delta + \nu_A(x)^\delta + \pi_A(x)^\delta = 1. \end{aligned}$$

Proofs of (ii) and (iii) are obvious.

(iv) Follows by noting that

$$f_{\alpha,1-\alpha}(A) = \left\{ \langle x, (\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + (1-\alpha)\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\} = d_\alpha(A).$$

(v) By (iv) we have $f_{0,1}(A) = d_0(A)$ and $f_{1,0}(A) = d_1(A)$. It follows by Theorem 4.1-(iii) that $f_{0,1}(A) = \overline{\Diamond A}$.

(vi) It follows by Theorem 4.1-(iv) that $f_{1,0}(A) = \overline{\Box A}$.

(viii) Since

$$f_{\beta,\alpha}(A) = \left\{ \langle x, (\nu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

and

$$f_{\alpha,\beta}(\bar{A}) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\nu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

we have

$$\overline{f_{\alpha,\beta}(\bar{A})} = \left\{ \langle x, (\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

and $\overline{f_{\alpha,\beta}(\bar{A})} = f_{\beta,\alpha}(A)$.

Since $\pi_{\Box A}(x) = 0$ and $\pi_{\Diamond A}(x) = 0$, the proof (vii), (x) and (xi) are obvious.

Theorem 4.3. For every $A \in \text{GIFS}_B$, and real numbers $\alpha, \beta, \gamma \in [0,1]$

- i. $g_{\alpha,\beta}(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow g_{\alpha,\beta}(A) \subset g_{\gamma,\beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow g_{\alpha,\beta}(A) \supset g_{\alpha,\gamma}(A)$,
- iv. $\tau \in [0,1] \Rightarrow g_{\alpha,\beta}(g_{\gamma,\tau}(A)) = \overline{g_{\gamma,\tau}(g_{\alpha,\beta}(\bar{A}))}$,
- v. $g_{\alpha,\beta}(C(A)) = I(g_{\alpha,\beta}(A))$,
- vi. $g_{\alpha,\beta}(I(A)) = C(g_{\alpha,\beta}(A))$,
- vii. $g_{\alpha,\beta}(\bar{A}) = g_{\beta,\alpha}(A)$,
- viii. $g_{1,1}(A) = \bar{A}$,
- ix. $g_{\alpha,\beta}(A \cap B) = g_{\alpha,\beta}(A) \cap g_{\alpha,\beta}(B)$,

$$x. \quad g_{\alpha,\beta}(A \cup B) = g_{\alpha,\beta}(A) \cup g_{\alpha,\beta}(B).$$

Proof. (i) Since

$$\begin{aligned} g_{\alpha,\beta}(A) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} v_A(x), \beta^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}, \\ \mu_{g_{\alpha,\beta}(A)}(x)^\delta + v_{g_{\alpha,\beta}(A)}(x)^\delta &= (\alpha^{\frac{1}{\delta}} v_A(x))^\delta + \left(\beta^{\frac{1}{\delta}} \mu_A(x) \right)^\delta, \\ &= \alpha v_A(x)^\delta + \beta \mu_A(x)^\delta, \\ &\leq v_A(x)^\delta + \mu_A(x)^\delta \leq 1. \end{aligned}$$

We have $g_{\alpha,\beta}(A) \in \text{GIFS}_B$.

(ii) Note that

$$g_{\alpha,\beta}(A) = \left\{ \langle x, \alpha^{\frac{1}{\delta}} v_A(x), \beta^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\},$$

and

$$g_{\gamma,\beta}(A) = \left\{ \langle x, \gamma^{\frac{1}{\delta}} v_A(x), \beta^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have $\alpha^{\frac{1}{\delta}} \leq \gamma^{\frac{1}{\delta}}$, $\alpha^{\frac{1}{\delta}} v_A(x) \leq \gamma^{\frac{1}{\delta}} v_A(x)$ and so $g_{\alpha,\beta}(A) \subset g_{\gamma,\beta}(A)$.

The proof of (iii) is similar to that of (ii).

(iv) Since

$$\begin{aligned} g_{\gamma,\tau}(A) &= \left\{ \langle x, \gamma^{\frac{1}{\delta}} v_A(x), \tau^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}, \\ g_{\alpha,\beta}(g_{\gamma,\tau}(A)) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} \tau^{\frac{1}{\delta}} \mu_A(x), \beta^{\frac{1}{\delta}} \gamma^{\frac{1}{\delta}} v_A(x) \rangle : x \in X \right\}, \\ &= \left\{ \langle x, (\alpha\tau)^{\frac{1}{\delta}} \mu_A(x), (\beta\gamma)^{\frac{1}{\delta}} v_A(x) \rangle : x \in X \right\}, \\ &= \overline{g_{\beta\gamma,\alpha\tau}(A)}, \end{aligned}$$

and

$$\begin{aligned} g_{\alpha,\beta}(\bar{A}) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} \mu_A(x), \beta^{\frac{1}{\delta}} v_A(x) \rangle : x \in X \right\}, \\ g_{\gamma,\tau}(g_{\alpha,\beta}(\bar{A})) &= \left\{ \langle x, \gamma^{\frac{1}{\delta}} \beta^{\frac{1}{\delta}} v_A(x), \tau^{\frac{1}{\delta}} \alpha^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}, \\ &= \left\{ \langle x, (\gamma\beta)^{\frac{1}{\delta}} v_A(x), (\alpha\tau)^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}, \\ &= g_{\gamma\beta,\alpha\tau}(A), \end{aligned}$$

we have $g_{\alpha,\beta}(g_{\gamma,\tau}(A)) = \overline{g_{\gamma,\tau}(g_{\alpha,\beta}(\bar{A}))}$.

(v) Follows by noting that

$$C(A) = \{ \langle x, \max_{y \in X} \mu_A(y), \min_{y \in X} \nu_A(y) \rangle : x \in X \},$$

and

$$\begin{aligned} g_{\alpha, \beta}(C(A)) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} \min_{y \in X} \nu_A(y), \beta^{\frac{1}{\delta}} \max_{y \in X} \mu_A(y) \rangle : x \in X \right\}, \\ &= \left\{ \langle x, \min_{y \in X} \alpha^{\frac{1}{\delta}} \nu_A(y), \max_{y \in X} \beta^{\frac{1}{\delta}} \mu_A(y) \rangle : x \in X \right\}, \\ &= I(g_{\alpha, \beta}(A)). \end{aligned}$$

(vi) Follows by noting that

$$I(A) = \{ \langle x, \min_{y \in X} \mu_A(y), \max_{y \in X} \nu_A(y) \rangle : x \in X \},$$

and

$$\begin{aligned} g_{\alpha, \beta}(I(A)) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} \max_{y \in X} \nu_A(y), \beta^{\frac{1}{\delta}} \min_{y \in X} \mu_A(y) \rangle : x \in X \right\}, \\ &= \left\{ \langle x, \max_{y \in X} \alpha^{\frac{1}{\delta}} \nu_A(y), \min_{y \in X} \beta^{\frac{1}{\delta}} \mu_A(y) \rangle : x \in X \right\}, \\ &= C(g_{\alpha, \beta}(A)) \quad , \quad \alpha, \beta \in [0, 1]. \end{aligned}$$

(vii) Let $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$ be a GIFSB. Then

$$\begin{aligned} \bar{A} &= \{ \langle x, \nu_A(x), \mu_A(x) \rangle : x \in X \}, \\ g_{\beta, \alpha}(A) &= \left\{ \langle x, \beta^{\frac{1}{\delta}} \nu_A(x), \alpha^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\}, \\ g_{\alpha, \beta}(\bar{A}) &= \left\{ \langle x, \alpha^{\frac{1}{\delta}} \mu_A(x), \beta^{\frac{1}{\delta}} \nu_A(x) \rangle : x \in X \right\}, \end{aligned}$$

and

$$\overline{g_{\alpha, \beta}(\bar{A})} = \left\{ \langle x, \beta^{\frac{1}{\delta}} \nu_A(x), \alpha^{\frac{1}{\delta}} \mu_A(x) \rangle : x \in X \right\},$$

so $\overline{g_{\alpha, \beta}(\bar{A})} = g_{\beta, \alpha}(A)$. Proof of (viii), (x) and (xi) are obvious.

Theorem 4.4. For every $A \in \text{GIFSB}$, and $\alpha, \beta, \gamma \in [0, 1]$

- i. $J_{\alpha, \beta}(A) \in \text{GIFSB}$,
- ii. $\alpha \leq \gamma \Rightarrow J_{\alpha, \beta}(A) \subset J_{\gamma, \beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow J_{\alpha, \beta}(A) \supset J_{\alpha, \gamma}(A)$,
- iv. $\diamond A = J_{1, 1}(A)$,
- v. $A = J_{0, 1}(A)$.

Proof. (i) Since

$$J_{\alpha, \beta}(A) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha \pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta \nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

and

$$\begin{aligned} \mu_{J_{\alpha,\beta}(A)}(x)^\delta + \nu_{J_{\alpha,\beta}(A)}(x)^\delta &= ((\mu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}})^\delta + \left(\beta^{\frac{1}{\delta}} \nu_A(x)\right)^\delta, \\ &= (\mu_A(x)^\delta + \alpha\pi_A(x)^\delta) + \beta\nu_A(x)^\delta, \\ &\leq \mu_A(x)^\delta + \pi_A(x)^\delta + \nu_A(x)^\delta = 1, \end{aligned}$$

we have $J_{\alpha,\beta}(A) \in \text{GIFS}_B$.

(ii) Note that

$$J_{\alpha,\beta}(A) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta\nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

and

$$J_{\gamma,\beta}(A) = \left\{ \langle x, (\mu_A(x)^\delta + \gamma\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta\nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have $(\mu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}} \leq (\mu_A(x)^\delta + \gamma\pi_A(x)^\delta)^{\frac{1}{\delta}}$ and so $J_{\alpha,\beta}(A) \subset J_{\gamma,\beta}(A)$.

The proof of (iii) is similar to that (ii). Proofs of (iv) and (v) are obvious.

Theorem 4.5. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $j_{\alpha,\beta}(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow j_{\alpha,\beta}(A) \subset j_{\gamma,\beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow j_{\alpha,\beta}(A) \supset j_{\alpha,\gamma}(A)$,
- iv. $j_{\alpha,\beta}(\bar{A}) = J_{\alpha,\beta}(A)$,
- v. $\overline{j_{1,1}(A)} = j_{1,1}(A)$,
- vi. $\bar{A} = j_{0,1}(A)$.

Proof. (i) Since

$$j_{\alpha,\beta}(A) = \left\{ \langle x, (\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta\mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

and

$$\begin{aligned} \mu_{j_{\alpha,\beta}(A)}(x)^\delta + \nu_{j_{\alpha,\beta}(A)}(x)^\delta &= ((\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}})^\delta + \left(\beta^{\frac{1}{\delta}} \mu_A(x)\right)^\delta, \\ &= (\nu_A(x)^\delta + \alpha\pi_A(x)^\delta) + \beta\mu_A(x)^\delta, \\ &\leq \mu_A(x)^\delta + \pi_A(x)^\delta + \nu_A(x)^\delta = 1. \end{aligned}$$

Finally, it can be concluded that $j_{\alpha,\beta}(A) \in \text{GIFS}_B$.

(ii) Note that

$$j_{\alpha,\beta}(A) = \left\{ \langle x, (\nu_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta\mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

and

$$j_{\gamma,\beta}(A) = \left\{ \langle x, (v_A(x)^\delta + \gamma\pi_A(x)^\delta)^{\frac{1}{\delta}}, (\beta\mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have $(v_A(x)^\delta + \alpha\pi_A(x)^\delta)^{\frac{1}{\delta}} \leq (v_A(x)^\delta + \gamma\pi_A(x)^\delta)^{\frac{1}{\delta}}$ and so $j_{\alpha,\beta}(A) \subset j_{\gamma,\beta}(A)$. The proof of (iii) is similar to that (ii). Proofs of (iv), (v) and (vi) are obvious.

Theorem 4.6. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $H_{\alpha,\beta}(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow H_{\alpha,\beta}(A) \subset H_{\gamma,\beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow H_{\alpha,\beta}(A) \supset H_{\alpha,\gamma}(A)$,
- iv. $H_{1,0}(A) = A$,
- v. $H_{1,1}(A) = A$.

Proof. (i) Since

$$H_{\alpha,\beta}(A) = \left\{ \langle x, (\alpha\mu_A(x)^\delta)^{\frac{1}{\delta}}, (v_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

and

$$\begin{aligned} \mu_{H_{\alpha,\beta}(A)}(x)^\delta + v_{H_{\alpha,\beta}(A)}(x)^\delta &= ((\alpha\mu_A(x)^\delta)^{\frac{1}{\delta}})^\delta + \left((v_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \right)^\delta, \\ &= \alpha\mu_A(x)^\delta + v_A(x)^\delta + \beta\pi_A(x)^\delta, \\ &\leq \mu_A(x)^\delta + \pi_A(x)^\delta + v_A(x)^\delta = 1, \end{aligned}$$

we have $H_{\alpha,\beta}(A) \in \text{GIFS}_B$.

(ii) Note that

$$H_{\alpha,\beta}(A) = \left\{ \langle x, (\alpha\mu_A(x)^\delta)^{\frac{1}{\delta}}, (v_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\},$$

and

$$H_{\gamma,\beta}(A) = \left\{ \langle x, (\gamma\mu_A(x)^\delta)^{\frac{1}{\delta}}, (v_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have $(\alpha\mu_A(x)^\delta)^{\frac{1}{\delta}} \leq (\gamma\mu_A(x)^\delta)^{\frac{1}{\delta}}$ and so $H_{\alpha,\beta}(A) \subset H_{\gamma,\beta}(A)$. The proof of (iii) is similar to that (ii). Proofs of (iv) and (v) are obvious.

Theorem 4.7. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $h_{\alpha,\beta}(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow h_{\alpha,\beta}(A) \subset h_{\gamma,\beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow h_{\alpha,\beta}(A) \supset h_{\alpha,\gamma}(A)$,
- iv. $h_{\alpha,\beta}(\bar{A}) = H_{\alpha,\beta}(A)$,
- v. $h_{1,0}(A) = \bar{A}$,
- vi. $h_{1,1}(A) = \overline{\delta A}$.

Proof. (i) Since

$$\begin{aligned}\mu_{h_{\alpha,\beta}(A)}(x)^\delta + \nu_{h_{\alpha,\beta}(A)}(x)^\delta &= (\alpha\nu_A(x)^\delta)^{\frac{1}{\delta}})^\delta + \left((\mu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \right)^\delta, \\ &= \alpha\nu_A(x)^\delta + \mu_A(x)^\delta + \beta\pi_A(x)^\delta, \\ &\leq \nu_A(x)^\delta + \mu_A(x)^\delta + \pi_A(x)^\delta = 1,\end{aligned}$$

we have $h_{\alpha,\beta}(A) \in \text{GIFS}_B$.

(ii) Note that

$$h_{\alpha,\beta}(A) = \left\{ \langle x, (\alpha\nu_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

and

$$h_{\gamma,\beta}(A) = \left\{ \langle x, (\gamma\nu_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta\pi_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have $(\alpha\nu_A(x)^\delta)^{\frac{1}{\delta}} \leq (\gamma\nu_A(x)^\delta)^{\frac{1}{\delta}}$ and so $h_{\alpha,\beta}(A) \subset h_{\gamma,\beta}(A)$.

The proof of (iii) is similar to that (ii). The proofs of (iv), (v) and (vi) are obvious.

Definition 4.2. Letting $\alpha, \beta \in [0,1]$ and $A \in \text{GIFS}_B$, we define the some operators of as follows

- i. $J_{\alpha,\beta}^*(A) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta\nu_A(x)^\delta))^{\frac{1}{\delta}}, (\beta\nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}$,
- ii. $J_{\alpha,\beta}^*(A) = \left\{ \langle x, (\nu_A(x)^\delta + \alpha(1 - \beta\mu_A(x)^\delta - \nu_A(x)^\delta))^{\frac{1}{\delta}}, (\beta\mu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}$,
- iii. $H_{\alpha,\beta}^*(A) = \left\{ \langle x, (\alpha\mu_A(x)^\delta)^{\frac{1}{\delta}}, (\nu_A(x)^\delta + \beta(1 - \alpha\mu_A(x)^\delta - \nu_A(x)^\delta))^{\frac{1}{\delta}} \rangle : x \in X \right\}$,
- iv. $h_{\alpha,\beta}^*(A) = \left\{ \langle x, (\alpha\nu_A(x)^\delta)^{\frac{1}{\delta}}, (\mu_A(x)^\delta + \beta(1 - \mu_A(x)^\delta - \alpha\nu_A(x)^\delta))^{\frac{1}{\delta}} \rangle : x \in X \right\}$.

Theorem 4.8. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $J_{\alpha,\beta}^*(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow J_{\alpha,\beta}^*(A) \subset J_{\gamma,\beta}(A)$,
- iii. $\beta \leq \gamma \Rightarrow J_{\alpha,\beta}^*(A) \supset J_{\alpha,\gamma}^*(A)$,
- iv. $J_{1,1}^*(A) = \emptyset$, A ,
- v. $J_{0,1}^*(A) = A$.

Proof. (i) Since

$$J_{\alpha,\beta}^*(A) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta\nu_A(x)^\delta))^{\frac{1}{\delta}}, (\beta\nu_A(x)^\delta)^{\frac{1}{\delta}} \rangle : x \in X \right\}.$$

and

$$\begin{aligned}\mu_{J_{\alpha,\beta}^*(A)}(x)^\delta + \nu_{J_{\alpha,\beta}^*(A)}(x)^\delta &= ((\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta\nu_A(x)^\delta))^{\frac{1}{\delta}})^\delta + \left(\beta^{\frac{1}{\delta}} \nu_A(x)^\delta \right)^\delta, \\ &= (\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta\nu_A(x)^\delta) + \beta\nu_A(x)^\delta), \\ &= ((1 - \alpha)\mu_A(x)^\delta + \beta(1 - \alpha)(\nu_A(x)^\delta) + \alpha),\end{aligned}$$

$$\leq ((1 - \alpha)\mu_A(x)^\delta + (1 - \alpha)(v_A(x)^\delta) + \alpha \leq 1,$$

we have $J_{\alpha,\beta}^*(A) \in \text{GIFS}_B$.

(ii) Note that

$$J_{\alpha,\beta}^*(A) = \left\{ \langle x, (\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta v_A(x)^\delta))^\frac{1}{\delta}, (\beta v_A(x)^\delta)^\frac{1}{\delta} \rangle : x \in X \right\},$$

and

$$J_{\gamma,\beta}^*(A) = \left\{ \langle x, (\mu_A(x)^\delta + \gamma(1 - \mu_A(x)^\delta - \beta v_A(x)^\delta))^\frac{1}{\delta}, (\beta v_A(x)^\delta)^\frac{1}{\delta} \rangle : x \in X \right\}.$$

Since $\alpha \leq \gamma$ we have

$$(\mu_A(x)^\delta + \alpha(1 - \mu_A(x)^\delta - \beta v_A(x)^\delta))^\frac{1}{\delta} \leq (\mu_A(x)^\delta + \gamma(1 - \mu_A(x)^\delta - \beta v_A(x)^\delta))^\frac{1}{\delta},$$

and so $J_{\alpha,\beta}^*(A) \subset J_{\gamma,\beta}^*(A)$.

The proof of (iii) is similar to that (ii). The proofs of (iv) and (v) are obvious.

Theorem 4.9. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $J_{\alpha,\beta}^*(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow J_{\alpha,\beta}^*(A) \subset J_{\gamma,\beta}^*(A)$,
- iii. $\beta \leq \gamma \Rightarrow J_{\alpha,\beta}^*(A) \supset J_{\alpha,\gamma}^*(A)$,
- iv. $J_{\alpha,\beta}^*(\bar{A}) = J_{\alpha,\beta}^*(A)$,
- v. $\overline{j_{1,1}^*(A)} = j_{1,1}^*(A)$,
- vi. $\bar{A} = j_{0,1}^*(A)$.

Proof. (i) Since

$$J_{\alpha,\beta}^*(A) = \left\{ \langle x, (v_A(x)^\delta + \alpha(1 - \beta\mu_A(x)^\delta - v_A(x)^\delta))^\frac{1}{\delta}, (\beta\mu_A(x)^\delta)^\frac{1}{\delta} \rangle : x \in X \right\},$$

and

$$\begin{aligned} \mu_{j_{\alpha,\beta}^*(A)}^*(x)^\delta + v_{j_{\alpha,\beta}^*(A)}^*(x)^\delta &= ((v_A(x)^\delta + \alpha(1 - \beta\mu_A(x)^\delta - v_A(x)^\delta))^\frac{1}{\delta})^\delta + \left(\beta^\frac{1}{\delta} \mu_A(x) \right)^\delta, \\ &= ((1 - \alpha)v_A(x)^\delta + (1 - \alpha)\beta\mu_A(x)^\delta) + \alpha, \\ &\leq (1 - \alpha)v_A(x)^\delta + (1 - \alpha)\mu_A(x)^\delta + \alpha \leq 1, \end{aligned}$$

we have $j_{\alpha,\beta}^*(A) \in \text{GIFS}_B$. The rest of the proof is similar to that of Theorem 4.8.

Theorem 4.10. For every $A \in \text{GIFS}_B$, and $\alpha, \beta, \gamma \in [0,1]$

- i. $H_{\alpha,\beta}^*(A) \in \text{GIFS}_B$,
- ii. $\alpha \leq \gamma \Rightarrow H_{\alpha,\beta}^*(A) \subset H_{\gamma,\beta}^*(A)$,
- iii. $\beta \leq \gamma \Rightarrow H_{\alpha,\beta}^*(A) \supset H_{\alpha,\gamma}^*(A)$,
- iv. $H_{1,0}^*(A) = A$,

$$v. \quad H_{1,1}^*(A) = \Box A.$$

The proof of Theorem 4.10 is similar to that of Theorem 4.8.

Theorem 4.11. For every $A \in \text{GIFS}_{\mathcal{B}}$, and for every three real numbers $\alpha, \beta, \gamma \in [0,1]$

- i. $h_{\alpha,\beta}^*(A) \in \text{GIFS}_{\mathcal{B}}$,
- ii. $\alpha \leq \gamma \Rightarrow h_{\alpha,\beta}^*(A) \subset h_{\gamma,\beta}^*(A)$,
- iii. $\beta \leq \gamma \Rightarrow h_{\alpha,\beta}^*(A) \supset h_{\alpha,\gamma}^*(A)$,
- iv. $h_{\alpha,\beta}^*(\bar{A}) = H_{\alpha,\beta}^*(A)$,
- v. $h_{1,0}^*(A) = \bar{A}$,
- vi. $h_{1,1}^*(A) = \Diamond \bar{A}$.

The proof of Theorem 4.11 is similar to that of Theorem 4.8.

5. CONCLUSIONS

We have introduced modal types of operators over $\text{GIFS}_{\mathcal{B}}$ and proved their relationships. An open problem is: definition of level operator, negation operator and other operators over $\text{GIFS}_{\mathcal{B}}$ and the study of their properties.

CONFLICTS OF INTEREST

The authors declare that they have no competing interests.

ACKNOWLEDGMENTS

The authors wish to thank the referee, for the careful reading of the paper and for the helpful suggestions and comments.

REFERENCES

- [1] Atanassov, K.T., "Intuitionistic fuzzy sets", Fuzzy Sets and Systems, 20(1): 87-96, (1986).
- [2] Atanassov, K.T., "Intuitionistic fuzzy sets", Springer Physica-Verlag, Heidelberg, (1999).
- [3] Atanassov, K.T., "More on intuitionistic fuzzy sets", Fuzzy Sets and Systems, 33: 37-45, (1989).
- [4] Atanassov, K.T., "On four intuitionistic fuzzy topological operators", Math Ware Soft Comput., 8: 65-70, (2001).
- [5] Atanassov, K.T., "On the modal operators deferent over the intuitionistic fuzzy set", Notes on Intuitionistic Fuzzy Sets, 10(1): 7-12, (2004).
- [6] Cornelis, C., Atanassov, K.T., Kerre, E., "Application of extended modal operators as a tool for constructing inclusion indicators over intuitionistic fuzzy sets", Soft Computing-Foundations and Theoretical Aspects., 133-142, (2004).
- [7] Atanassov, K.T., Dimitrov, D., "On the negations over intuitionistic fuzzy sets", Annual of "Informatics" Section Union of Scientists in Bulgaria, 1: 49-58, (2008).

- [8] Hinde, C.J., Atanassov, K.T., “On intuitionistic fuzzy negations and intuitionistic fuzzy modal operators with contradictory evidence”, 5th WSEAS / IASME International Conference on Engineering Education (EE'08), Heraklion, Greece, July 22-24, (2008).
- [9] Parvathi, R., Geetha, S.P., “A note on properties of temporal intuitionistic fuzzy sets”, Notes on Intuitionistic Fuzzy Sets, 15(1): 42–48, (2009).
- [10] Atanassov, K.T., “On the operators partially extending the extended intuitionistic fuzzy operators from modal type”, Notes on Intuitionistic Fuzzy Sets, 18(3): 16–22, (2012).
- [11] Srinivasan, R., Palaniappan, N., “Some operators on intuitionistic fuzzy sets of root type”, Annals of Fuzzy Mathematics and Informatics, 4(2): 377-383, (2012).
- [12] Sharma, P.K., “Modal operator $F_{\alpha,\beta}(A)$ in intuitionistic fuzzy groups”, Annals of Pure and Applied Mathematics, 7(1): 19-28, (2014).
- [13] Yilmaz, S., Cuvalcioglu, G., “On level operators for temporal intuitionistic fuzzy sets”, 18th Int. Conf. on IFSs, Sofia, 10–11 May 2014, Notes on Intuitionistic Fuzzy Sets, 20(2): 6–15, (2014).
- [14] Baloui Jamkhaneh, E., Nadarajah, S., “A new generalized intuitionistic fuzzy sets”, Hacettepe Journal of Mathematics and Statistics, 44 (6): 1537–1551, (2015).
- [15] Baloui Jamkhaneh, E., “The operators over the generalized intuitionistic fuzzy sets”, International Journal of Nonlinear Analysis and Applications, 8(1): 11-21, (2017).
- [16] Baloui Jamkhaneh, E., Nadi Ghara A., “Four new operators over generalized intuitionistic fuzzy sets”, Journal of New Theory, 18: 12-21, (2017).
- [17] Baloui Jamkhaneh, E., Garg, H., “Some new operations over the generalized intuitionistic fuzzy sets and their application to decision making process”, Granular Computing, DOI 10.1007/s41066-017-0059-0, (2017).