






The lipide soluble vitamin contents of some *Onobrychis* Miller (*Fabaceae*) taxa

Bazı *Onobrychis* Miller (*Fabaceae*) Taksonları'nın yağda çözünen vitamin içeriği

Irfan EMRE¹ , Hakan SEPET² , Murat KURSAT³ , Muammer BAHSI¹ , Okkes YILMAZ⁴ , Ahmet SAHİN⁵ 

¹Firat University, Faculty of Education, Department of Primary Education, Elazig, Turkey

²Kirsehir Ahi Evran University, Faculty of Engineering, Department of Environmental Engineering, Kirsehir, Turkey.

³Bitlis Eren University, Faculty of Science and Arts, Department of Biology, Bitlis, Turkey

⁴Firat University, Faculty of Science, Department of Biology, Elazig, Turkey.

⁵Erciyes University, Faculty of Education, Department of Secondary Science and Mathematics Education, Kayseri, Turkey

Eser Bilgisi / Article Info

Araştırma makalesi / Research article

DOI: 10.17474/artvinofd.555426

Sorumlu yazar / Corresponding author

Irfan EMRE

e-mail: iemre@firat.edu.tr

Geliş tarihi / Received

18.04.2019

Düzeltilme tarihi / Received in revised form

25.09.2019

Kabul Tarihi / Accepted

06.10.2019

Elektronik erişim / Online available

30.10.2019

Keywords:

Fabaceae

Onobrychis

Vitamin

Anahtar kelimeler:

Fabaceae

Onobrychis

Vitamin

Abstract

The goal of this study is to determine the lipid-soluble vitamin contents in seeds of the some *Onobrychis* Miller (*Fabaceae*) taxa by using HPLC. Samples were collected from the natural habitats. Studied materials were dissolved in acetonitrile/methanol (75/25 v/v) and were injected 50 µL to HPLC instrument (Shimadzu, Kyoto Japan). According to data obtained from present study showed that *O. hypargyrea*, *O. viciifolia*, *O. caput-galli*, *O. fallax* and *O. oxyodonta* var. *armena* have high lipide-soluble vitamin contents. Present study found that *O. oxyodonta* var. *armena* (1777.27±6.24 µg/g), *O. fallax* (916.0±4.51 µg/g) *O. hypargyrea* (809.7±5.03 µg/g) and *O. viciifolia* (399.7±3.54 µg/g) have highest beta-caroten content. Also, *O. caput-galli* has high beta caroten content (73.3±94 µg/g). on the other hand, it was found that *O. fallax* has highest gamma-tocopherol content (1401.2±8.76 µg/g). *O. viciifolia* (574.9±2.35 µg/g), *O. caput-galli* (410.1±4.56 µg/g), *O. oxyodonta* var. *armena* (267.7±3.68 µg/g), *O. podporea* (162.5±2.14 µg/g) were the other high gamma tocopherol content. Whereas, retinol, retinol acetate and r-tocopherol contents were found absent or trace amounts in the present study.

Özet

Bu çalışmanın amacı, bazı *Onobrychis* Miller (*Fabaceae*) taksonlarının tohumlarındaki yağda çözünen vitamin içeriğini HPLC kullanarak belirlemektir. Doğal yaşam alanlarından örnekler alındı. Çalışılan malzemeler asetonitril / metanol (75/25 h / h) içinde çözüldü ve HPLC cihazına (Shimadzu, Kyoto Japonya) 50 µL enjekte edildi. Bu çalışmadan elde edilen verilere göre *O. hypargyrea*, *O. viciifolia*, *O. caput-galli*, *O. fallax* ve *O. oxyodonta* var. *armena*'nın lipitte çözünen vitamin içeriğinin yüksek olduğunu göstermiştir. Bu çalışma *O. oxyodonta* var. *armena* (1777.27 ± 6.24 µg / g), *O. fallax* (916.0 ± 4.51 µg / g) *O. hypargyrea* (809.7 ± 5.03 µg / g) ve *O. viciifolia* (399.7 ± 3.54 µg / g) en yüksek beta karoten içeriğine sahiptir. Ayrıca, *O. caput-galli* de yüksek beta karoten içeriğine sahiptir (73.3 ± 94 µg / g). Öte yandan, *O. fallax*'ın en yüksek gamma-tokoferol içeriğine sahip olduğu belirlendi (1401.2 ± 8.76 µg / g). *O. viciifolia* (574.9 ± 2.35 µg / g), *O. caput-galli* (410.1 ± 4.56 µg / g), *O. oxyodonta* var. *armena* (267.7 ± 3.68 µg / g), *O. podporea* (162.5 ± 2.14 µg / g) diğer yüksek gama tokoferol içeriğine sahip taksonlardır. Diğer taraftan bu çalışmada retinol, retinol asetat ve r-tokoferol içerikleri bulunmamakta veya eser miktarda bulunmaktadır.

INTRODUCTION

Onobrychis Miller, is a member of the *Fabaceae*, includes about 170 perennial and annual species in two subgenera (Aktoklu 1995, Karamian et al. 2012, Avci et al. 2013). The genus distributed in Europe, Asia, North America and Africa (Yildiz et al. 1999, Pavlova and Monova 2000, Kaveh et al. 2019). Turkey is one of the most significant center of the genus and it is represented by 55 taxa which 28 of them are endemic (Duman and Vural 1990, Davis et al. 1988, Aktoklu 2001, Avci and Kaya 2013).

The members of *Onobrychis* Miller are important agricultural sources as a forage, fodder legume or ornamental (Ranjbar et al. 2010, Carbonero et al. 2011). The species of genus also used to improve the quality of the soil by serving fix atmospheric nitrogen and they contribute to the organic structure of soil with root systems (Ozaslan Parlak and Parlak 2008, Arslan and Ertugrul, 2010, Yildiz and Ekiz 2014). Biochemical studies performed *Onobrychis* Miller taxa showed that the genus have antioxidant, antibacterial and antifungal effects (Karakoca et al. 2015, Karamanian and Asadbegy, 2016,

Bektas et al. 2018). However, these biochemical studies of genus extremely limited. Therefore, it was aimed to contribute the such studies of *Onobrychis* Miller by determining the lipide-soluble vitamins in this study.

MATERIAL AND METHODS

Table 1. Localities of collected plant samples

Taxa	Section	Region	Locality	Altitude
<i>O. hypargyrea</i> Boiss.	<i>Hymenobryhis</i>	B2, Kutahya	Usak Gediz road Abide bridge locality	690 m
<i>O. viciifolia</i> Scop.	<i>Onobrychis</i>	B2, Usak	From Usak to Banaz 7th km	100m
<i>O. cappadocica</i> Boiss.	<i>Hymenobryhis</i>	B7, Elazig	Firat University Campus, Faculty of Engineering locality	1060 m
<i>O. podporea</i> Širj.	<i>Onobrychis</i>		Usak Gediz road 30. km	740m
<i>O. caput-galli</i> (L) Lam	<i>Lophobrychis</i>	B2, Manisa	3 km from Kula to Alasehir, Kula dam lake locality	731 m
<i>O. galegifolia</i> Boiss.	<i>Hymenobryhis</i>	B7, Elazig	Elazig-Harput road	1230m
<i>O. fallax</i> Freyn & Sint. ex Freyn var. <i>fallax</i>	<i>Onobrychis</i>	B7, Elazig	Firat University Campus, Faculty of Engineering locality	1060 m
<i>O. oxyodonta</i> Boiss.var. <i>armena</i> (Boiss. & Huet) Aktoklu	<i>Onobrychis</i>	B2, Usak	Usak between Akarca	972m

Extraction of plant materials

1 g seed used to analyse the lipide-soluble vitamin contents. The seeds are finely ground in a mill and were then extracted with hexane/isopropanol (3:2 v/v) (Hara and Radin, 1978). Extracts were centrifuged at 10.000 g for 5 minutes and filtered. The solvent was then removed on a rotary evaporator at 40°C. After that lipide-soluble vitamins were extracted based on the method of Sánchez-Machado (2002) with minor modifications. The experiment was repeated three times.

Chromatographic analysis and quantification of lipid-soluble vitamins

Seed materials were dissolved in acetonitrile/methanol (75/25 v/v) and were injected 50 µL to HPLC instrument (Shimadzu, Kyota Japan). Supelcosil TM LC18 (250 x 4.6 mm, 5 mm, Sigma, USA) was used as column. The mobile

Collection of plant materials

In the present study, lipid-soluble vitamin contents in mature seeds of the *Onobrychis* L. taxa were examined. Sample plants were gathered from the natural habitats and details about the materials are explained in table I.

phase was acetonitrile/methanol (75/25 v/v) and the elution was performed at a flow-rate of 1 ml/min. The temperature of analytical column was maintained at 40 °C. Detection was conducted at 320 nm for retinol (vitamin A) and retinol acetate, and 215 nm for δ-tocopherol, vitamin D2 and D3, α-tocopherol, α-tocopherol acetate, 235 nm for vitamin K1. Identification of the individual vitamins were performed by frequent comparison with authentic external standard mixtures analyzed under the same conditions. Class Vp 6.1 software assisted at workup of the data (Yilmaz et al. 2007). The results of analysis were expressed as µg/g for samples.

RESULTS

The lipide-soluble vitamin contents of studied *Onobrychis* species were given in table 2.

Table 2. The lipide-soluble vitamin contents of studied *Onobrychis* species

Taxa	Lipide-soluble vitamins (µg/g)									
	Beta carotene	Gamma tocopherol	R-tocopherol	D2	D3	a-tocopherol	a-tocopherol acetate	K1	Retinol	Retinol acetate
<i>O. hypargyrea</i>	809.7±5.03	33.3±.97	0.7±0.05	3.3±0.2	51.2±1.12	22.2±2.64	10.2±0.97	4.9±0.24	0.4±0.15	0.3±0.01
<i>O. viciifolia</i>	399.7±3.54	574.9±2.35	1.4±0.06	0.3±0.1	64.2±2.28	1.8±0.4	0.8±0.02	5.2±0.33	0.2±0.11	0.4±0.03
<i>O. cappadocia</i>	-	-	0.1±0.01	2.7±0.5	55.4±0.97	0.9±0.02	0.2±0.01	6.3±0.52	1.0±0.12	0.5±0.01
<i>O. podporea</i>	-	162.5±2.14	0.3±0.01	1.1±0.3	39.3±0.75	8.0±0.7	-	3.7±0.21	0.4±0.05	0.6±0.02
<i>O. caput-galli</i>	73.3±0.94	410.1±4.56	-	-	36.7±1.1	0.4±0.01	0.6±0.02	-	0.3±0.07	0.3±0.03
<i>O. galegifolia</i>	-	-	3.0±0.7	15.0±1.1	55.2±2.1	6.1±0.3	-	-	0.9±0.08	0.7±0.03
<i>O. fallax</i>	916.0±4.51	1401.2±8.76	-	1.8±0.08	66.7±1.2	2.0±0.05	1.9±0.74	1.8±0.13	0.8±0.03	0.9±0.03
<i>O. oxyodonta</i> <i>var. armena</i>	1777.2±6.24	267.7±3.68	-	-	58.3±1.04	2.0±0.04	0.6±0.1	1.6±0.1	0.4±0.06	0.6±0.01

It was found that *O. hypargyrea*, *O. viciifolia*, *O. caput-galli*, *O. fallax* and *O. oxyodonta var. armena* have high lipide-soluble vitamin content based on results of this study (table 2). Present study showed that *O. oxyodonta var. armena* (1777.27±6.24 µg/g), *O. fallax* (916.0±4.51 µg/g) *O. hypargyrea* (809.7±5.03 µg/g) and *O. viciifolia* (399.7±3.54 µg/g) have quite highest beta-caroten content. *O. caput-galli* has high beta caroten content (73.3±.94 µg/g). It was found that *O. fallax* has highest gamma-tocopherol content (1401.2±8.76 µg/g). In addition to, *O. viciifolia* (574.9±2.35 µg/g), *O. caput-galli* (410.1±4.56 µg/g), *O. oxyodonta var. armena* (267.7±3.68 µg/g), *O. podporea* (162.5±2.14 µg/g) high gamma-tocopherol content. Furthermore, *O. hypargyrea* has low gamma tocopherol content (33.3±.97µg/g) while *O. cappadocia* and *O. galegifolia* don't have gamma tocopherol content. Furthermore, present study showed that *O. taxa* have D3 vitamin content between 66.7±1.2 µg/g (*O. fallax*) and 36.7±1.1 µg/g (*O. caput-galli*). A-tocopherol content of studied *O. species* range from 0.4±0.01 µg/g (*O. caput-galli*) to 22.2±2.64 µg/g (*O. hypargyrea*). Also, *O. hypargyrea* has high a-tocopherol content 10.2±0.97 µg/g among studied *O. species*. Moreover, K1 content of *O. species* varied from 1.6±0.1 µg/g (*O. oxyodonta var. armena*) from 6.3±0.52 µg/g (*O. cappadocia*) except for *O. caput-galli* and *O. galegifolia* which don't have K1 content. Retinol and retinol acetate contents of *O. species* found lowest or trace amounts in the present study.

DISCUSSION

Legumes are consumed high levels especially Asia, Africa and South America (Frias et al. 2005) and studies showed that legumes have complex carbohydrates, vitamins, fibers, polyphenols (Obob 2006, Amarowicz and Pegg 2008). These bioactive compounds play significant role many diseases such as cancer, diabetes (Frias et al. 2005, Arslan, 2017). Lipide-soluble phytonutrients such as carotenoids and tocopherols have been reported to inhibit the risk of cardiovascular, cancer, eye pathologies and diabetes (Monge-Rajos and Campos 2011, Nadeau et al. 2013). Also, they have important roles in anti-inflammatory processes and immune system by scavenging cells against free radical damages (McDowell 2000, Chou et al. 2007, Fernandez-Marin et al. 2014).

Beta-carotene is considered to be pro-vitamin which has the ability to be converted into vitamin A (Hojer et al. 2012). Beta-carotene, is considered to be pro-vitamins because they have the ability to be converted into vitamins (vitamin A or retinol) by the animal (Hojer et al. 2012). On the other hand, vitamin E is, a lipophilic structure and major constituent of cell membrane (Kappus and Diplock 1992), externally intaken in foods or supplements because it isn't generating by humans (Berman and Brodaty 2004). Tocopherols have protective role against free radical damages in cells by interrupting the chain reactions (Bramley et al. 2000). Present study showed that some of studied *Onobrychis* species have highest beta-carotene and gamma-tocopherol contents. A study done Wyatt et al. (1998) showed that all of the

legumes analyzed showed the presence of γ -tocopherol in relatively high levels, with the exception of black beans. Fernandez-Marin et al. (2014) found that of all tocopherols, γ -tocopherol was the most abundant isoform in all species, apart from *Vigna* and *Arachis*, where δ -tocopherol and α -tocopherol were the main isoforms, respectively. Also, they found that total carotenoids were between 0.9 ± 0.2 $\mu\text{g/g}$ and 17.7 ± 2.2 $\mu\text{g/g}$ (Fernandez-Marin et al. 2014). Another study done by Boschini and Arnoldi (2011) showed that legume seeds have 0.3-2.99 mg/100 g tocopherol content. It was reported that legumes have contain only γ -tocopherols (86.1–146.8 mg/kg) study done by Cho et al. (2007). Also, Cho et al. (2007) determined the carotene content of legumes is 9.2 ± 10 mg/kg. El-Qudah (2014) identified legumes including *Vicia*, *Lens*, *Phaseolus* and *Cicer* have appreciable amounts of carotenoid. However, Mamatha et al. (2011) found that studied legumes including *Phaseolus*, *Vigna*, *Lens* and *Cicer* have lowest a-and b-carotene contents. A-tocopherol content of *O.* was found between 22.2 ± 2.64 $\mu\text{g/g}$ and 1.8 ± 0.4 $\mu\text{g/g}$ while K1 content of *O.* was found between 1.6 ± 0.1 $\mu\text{g/g}$ and 10.2 ± 0.97 $\mu\text{g/g}$ (except for *O. caput-galli* and *O. galegifolia* which don't have K1 content) in present study. Arslan (2017) indicated that legumes include K vitamin together with vitamin B1, B2, B6, vitamin C, vitamin E. Furthermore, it was found that studied *O.* species have high D3 content (66.7 ± 1.2 - 36.7 ± 1.1 $\mu\text{g/g}$) in this study. Sahin et al. (2009) found that *Lathyrus* taxa, the other genus of legumes, have high vitamin D3. Also, they determined that *Lathyrus* has high δ -tocopherol, α -tocopherol, α -tocopherol acetate contents (Sahin et al. 2009). On the other hand, present work demonstrated that r-tocopherol, retinol, retinol acetat, vitamin D2 (except for *O. galegifolia*) contents of *O.* has lowest. Similarly, Sahin et al. (2009) found that retinol, retinol acetate, vitamin D2 were trace amounts in their work.

REFERENCES

- Aktoklu E (1995) Türkiye'de yetişen *Onobrychis* Miller (Fabaceae) türlerinin revizyonu, İnönü Üniversitesi Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Doktora tezi, 135 s.
- Aktoklu E (2001) Two New Varieties and a New Record in *Onobrychis* from Turkey. *Turk J Bot.* 25 (5):359-363.

- Amarowicz R, Peg, RB (2008) Legumes as a source of natural antioxidants. *Eur. J. Lipid Sci. Technol.* 110 (10): 865–878.
- Arslan E, Ertugrul K (2010) Genetic relationships of the genera *Onobrychis*, *Hedysarum*, and *Sartoria* using seed storage proteins. *Turk J Biol* 34(1): 67-73.
- Arslan M (2017) Diversity for vitamin and amino acid content in grass pea (*Lathyrussativus* L.). *Legume Research* 40 (5): 803-810.
- Avci S, Kaya MD (2013) Seed and germination characteristics of wild *O.* taxa in Turkey. *Turk J Agric For.* 37 (5): 555-560
- Bektas E, Kaltalioglu K, Sahin H, Turkmen Z, Kandemir A (2018) Analysis of phenolic compounds, antioxidant and antimicrobial properties of some endemic medicinal plants. *International Journal of Secondary Metabolite* 5 (2): 75–86.
- Berman K, Brodaty H (2004) Tocopherol (Vitamin E) in Alzheimer's Disease and Other Neurodegenerative Disorders. *CNS Drugs* 18 (12): 807-825.
- Boschin G, Arnoldi A (2011) Legumes are valuable sources of tocopherols. *Food Chemistry* 127 (3): 1199–1203.
- Bramley PM, Elmadfa I, Kafatos A, Kelly FJ, Manios Y, Roxborough HE (2000) Vitamin E (review). *Journal of Science and Food Agriculture* 80(7): 913–938.
- Carbonero CH, Mueller-Harvey I, Brown TA, Smith L (2011). Sainfoin (*O. viciifolia*): a beneficial forage legume. *Plant Genetic Resources: Characterization and Utilization* 9(1): 70–85.
- Cho YS, Yeum KJ, Chen CY, Beretta G, Tang G, Krinsky NI, Yoon S, Lee-Kim YC, Blumberg JB, Russell RM (2007) Phytonutrients affecting hydrophilic and lipophilic antioxidant activities in fruits, vegetables and legumes. *Journal of the Science of Food and Agriculture* 87(5):1096 –1107.
- Davis PH (1988) *Flora of Turkey and The East Aegean Island*. Edinburgh University Press. No:10, Edinburgh.
- Duman H, Vural M (1990) New taxa from south Anatolia. I. *Turk J Bot* 14(1):45-48.
- El-Qudah JM (2014) Estimation of Carotenoid Contents of Selected Mediterranean Legumes by HPLC. *World Journal of Medical Sciences* 10 (1): 89-93.
- Fernandez-Marin B, Milla R, Martin-Robles N, Arc E, Kranner I, Becerril JM, Garcia-Plazaola I (2014) Side-effects of domestication: cultivated legume seeds contain similar tocopherols and fatty acids but less carotenoids than their wild counterparts. *BMC Plant Biology* 14 (1599): 1-11.
- Frias MJ, Miranda ML, Doblado R, Vidal-Valverde C (2005) Effect of germination and fermentation on the antioxidant vitamin content and antioxidant capacity of *Lupinus albus* L. var. *multolupa*. *Food Chemistry* 92(2): 211–220.
- Hara A, Radin NS (1978). Lipid extraction of tissues with a low-toxicity solvent. *Anal.Biochem.* 90 (1): 420-426.
- Hojer A, Adler S, Martinsson K, Jnesen SK, Steinshamn H, Thuen E, Gustavsson AM (2012) Effect of legume–grass silages and a-tocopherol supplementation on fatty acid composition and a-tocopherol, b-carotene and retinol concentrations in organically produced bovine milk. *Livestock Science* 148(3): 268–281.
- Kappus H, Diplock AT (1992) Tolerance and safety of vitamin E: a toxicological position report. *Free Radic Biol Med* 13(1): 55–74.

- Karakoca K, Asan-Ozusaglam M, Cakmak YS, Teksen M (2015) Phenolic compounds, biological and antioxidant activities of *O. armena* Boiss. & Huet flower and root extracts. *Chiang Mai University Journal of Natural of Sciences* 42(2): 376–392.
- Karamian R, Asadbegy M (2016) Antioxidant activity, total phenolic and flavonoid contents of three *Onobrychis* species from Iran. *Journal of Pharmaceutical Sciences* 22(2): 112–119.
- Kaveh A, Kazempour-Osaloo S, Amirahmadi A, Maassoumi A, Schneeweiss GM (2019) Systematics of *O. sect. Heliobrychis* (Fabaceae): morphology and molecular phylogeny. *Plant Systematics and Evolution* 305(1): 33–48.
- McDowell LR (2000) Vitamins in animal and human nutrition. Iowa State University Press: Ames, Iowa, USA.
- Mamatha BS, Sageetha RK, Baskaran V (2011) Provitamin-A and xanthophyll carotenoids in vegetables and food grains of nutritional and medicinal importance. *International Journal of Food Science and Technology* 46(4): 315–323.
- Monge-Rojas R, Campos H (2011) Tocopherol and carotenoid content of foods commonly consumed in Costa Rica. *Journal of Food Composition and Analysis* 24(2): 202–216.
- Nadeau E, Lindqvist H, Jnesen SK, Nilsdotter N, Gustavsson A (2013) Variations in α -tocopherol and β -carotene concentrations in forage legumes and grasses harvested at different sites and maturity stages. In: *Proceedings of the 22nd International Grassland Congress*. pp 643-646.
- Oboh G (2006) Antioxidant properties of some commonly consumed and under-utilized tropical legumes. *European Food Research Technology* 224(1): 61–65.
- Ozaslan Parlak A, Parlak M (2008) Effect of salinity in irrigation water on some plant development parameters of sainfoin (*O. viciifolia* Scop.) and soil salinity. *Tarım Bilim Dergisi* 14(4): 320–325.
- Pavlova DK, Manova VI (2000) Pollen morphology of the genera *O.* and *Hedysarum* (Hedysarea, Fabaceae) in Bulgaria. *Ann Bot Fenn* 37(3): 207–217.
- Ranjbar M, Karamian VE (2010) *O. bakuensis* (Fabaceae), a new species from Azerbaijan. *Ann. Bot. Fennici* 47(3): 233-236.
- Sahin A, Emre I, Yilmaz O, Genc H, Karatepe M (2009) Vitamin and fatty acid contents in seeds of some taxa belonging to genus *Lathyrus* L. growing in Turkey. *Acta Botanica Gallica* 156 (3): 331-339.
- Sánchez-Machado DI, Lopez-Hernandez J, Paseiro-Losado P (2002) High-performance liquid chromatographic determination of α -tocopherol in macroalgae. *Journal of Chromatography A* 976(1): 277–284.
- Wyatt CJ, Carballido SP, Mendez RO (1998) α - and γ -Tocopherol Content of Selected Foods in the Mexican Diet: Effect of Cooking Losses. *J. Agric. Food Chem.* 46(11): 4657–4661.
- Yilmaz O, Keser S, Tuzcu M, Cetintas B (2007) Resveratrol (trans-3,4',5-trihydroxystilbene) decreases lipid peroxidation level and protects antioxidant capacity in sera and erythrocytes of old female Wistar rats induced by the kidney carcinogen potassium bromate. *Envir. Toxicol. Pharmacol.* 24(2): 79-85.
- Yildiz B, Ciplak B, Aktoklu E (1999) Preliminary phylogeny of sections of genus *Onobrychis* Miller (Fabaceae) with references of fruit morphology. *Isr. J. Plant Sci.* 47(4): 269-282.
- Yildiz M, Ekiz H (2014) The effect of sodium hypochlorite solutions on in vitro seedling growth and regeneration capacity of sainfoin (*O. viciifolia* Scop.) hypocotyl explants. *Can. J. Plant Sci.* 94(7): 1161-1164.